



TITLE:

On Homology 3-Spheres Which Bound Contractible 4-Manifolds (低 次元多様体の構造と分類について)

AUTHOR(S):

MARUYAMA, NORIKO

CITATION:

MARUYAMA, NORIKO. On Homology 3-Spheres Which Bound Contractible 4-Manifolds (低次元多様体の構造と分類について). 数理解析研究所講究録 1981, 417: 122-129

ISSUE DATE:

1981-02

URL:

<http://hdl.handle.net/2433/102485>

RIGHT:

On homology 3-spheres
which bound contractible 4-manifolds

NORIKO MARUYAMA

TSUDA COLLEGE

§1. Introduction.

In this note, we consider the question which homology 3-spheres bound contractible (or acyclic) 4-manifolds. This question is important in the problem of classifying homology 3-spheres. Some families of homology 3-spheres which do bound contractible 4-manifolds are given by Akbulut and Kirby and others in [A-K], [C-H] and [M]. All known examples are the Breiskorn homology 3-spheres.

We deal with the homology 3-spheres of plumbing type. A closed 3-manifold is of plumbing type if it is the boundary of a regular neighbourhood of smooth, normally immersed surface in an oriented 4-manifold. The Breiskorn homology 3-spheres are of course of plumbing type, but all of homology 3-spheres of plumbing type is not a Breiskorn homology 3-sphere.

Throughout this note we use the terminology of Kirby's calculus on framed links. Since Kirby's calculus is widely known, we omit the minute details. We refer to [K] for the fundamentals of Kirby's calculus. Every time we draw a framed link in S^3 , we mean the corresponding 4-manifold obtained by attaching 2-handles on the boundary of B^4 along the framed link. In this note

\approx means a diffeomorphism and \cong means a diffeomorphism between the boundaries of manifolds. "Blowing up" means taking connected sum with CP^2 or $-CP^2$. "Blowing down" means replacing the neighbourhood of an embedded 2-sphere ($=$ the D^2 -bundle over S^2) with B^4 when the Euler characteristic of the bundle is ± 1 .

In §2 we define a contractible 4-manifold by generalizing so called Mazur manifold. Applying Kirby's calculus on framed links in §3, we sketch the proof of the following result:

Theorem 1. ([M1], [M2])

A. A Brieskorn homology 3-spheres $\Sigma(2n+1, 2n+2, 2n+3)$ bounds a contractible 4-manifold.

B. A homology 3-sphere $\Sigma(n)$ of plumbing type given by the following framed link $\mathcal{V}(n)$ ($n \geq 1$) bounds a contractible 4-manifold where


$$\Sigma(n) = \partial \left(\mathcal{V}(n): \begin{array}{c} \begin{array}{c} \text{Top row of circles:} \\ \text{Left: } -2, -1, -2, -2, \dots, -2, -2 \\ \text{Middle: } -4, \dots, n \end{array} \\ \begin{array}{c} \text{Bottom row of circles:} \\ \text{Left: } -2n-3, -2, -2, \dots, -2, -2 \\ \text{Middle: } -2, \dots, n-1 \end{array} \end{array} \right).$$

The statement A is stronger than the result of Martin in [M]: $\Sigma(2n+1, 2n+2, 2n+3)$ bounds an acyclic 4-manifold. But it is the special case of the results of Casson and Harer [C-H]. ~~We note about~~

the family of homology 3-spheres in the statement B of Theorem 1 that $\Sigma(1)$ is the Breiskorn homology 3-sphere $\Sigma(2, 5, 7)$ and that $\Sigma(n)$ ($n \geq 2$) is not a Breiskorn homology 3-sphere [M2].

Another method to give an example of homology 3-sphere which bounds a contractible 4-manifold is to perform ± 1 -surgery on S^3 along a slice knot in S^3 [G]. By Kirby's calculus we can show

Theorem 2. ([M1]) A Breiskorn homology 3-sphere $\Sigma(2, 3, 6n \pm 1)$ is obtained by a ± 1 -surgery on a knot $K^\pm(n)$ ($n \geq 1$) where




Since we can show only $K^+(2)$ is a slice knot among $K^\pm(n)$'s ($n \geq 1$), we restate that

Corollary. $\Sigma(2, 3, 13)$ bounds a contractible 4-manifold.

§2. Construction of a contractible 4-manifold.

$W^4 = W(n+1, n; r)$ is defined as a 4-manifold obtained by adding two 2-handles to B^4 along the following framed link of two components in $S^3 = \partial B^4$ and surgering S^2 corresponding to the 0-framed unknotted circle with a dot:

$W(n+1, n; r):$



where the r -framed circle α links the 0 -framed circle with a dot algebraically once.

By the definition W is a 4 -manifold with its handle decomposition consisting one 1 -handle and one 2 -handle attached along the r -framed circle α lies in $S^1 \times S^2 = \partial(B^4 \times H^1) = \partial(S^1 \times B^3)$ being homotopic but not isotopic to the standard embedding $S^1 \times (*) \subset S^1 \times S^2$. That is the 2 -handle cancels the 1 -handle homotopically. Therefore W can be seen as a generalized contractible Mazur 4 -manifold with boundary.

The following theorem gives a description of the boundaries of some of these contractible 4 -manifolds of Mazur types.

Theorem 1.


A. $\Sigma(2n+1, 2n+2, 2n+3) \approx \partial W(n+1, n; 2n+2) \quad (n \geq 1).$

B. $\Sigma(n) \approx \partial W(n+1, n; 2n+1) \quad (n \geq 1).$


§3. Proof.

For want of space we prove only the statement of Theorem 1.B. We refer to [M1] and [M2] for the explicit proofs of the results.


Proof of Theorem 1.B. We show that $\mathcal{V}(n)$ can be deformed by Kirby's calculus into the framed link defining $W(n+1, n; 2n+1)$.




\approx
Blow down



\approx
Blow down




=





\approx
Handle adding
corresponding to

$$\begin{cases} h_1 \rightarrow h_1 \\ h_2 \rightarrow h_2 + h_1 \end{cases}$$



\approx

Blow down the -1
framed circles $n+1$ times
one after the other


 $=$

 \approx
 $W(n+1, n; 2n+1).$

Surger S^2
corresponding to



This completes the proof of Theorem 1. B.

Concluding remarks

(i) From a different point of view we can show that the Breiskorn homology 3-sphere $\Sigma(2, 7, 19)$ bounds a contractible 4-manifold, applying Kirby's calculus.

(ii) The proof of (i) and related topics will be described in another paper.

References

- [A-K] S. Akbult and R. C. Kirby, Mazur manifolds. (preprint)
- [C-H] A. J. Casson and J. L. Harer, Some Breiskorn homology 3-spheres which bound contractible 4-manifolds. (preprint)
- [G] C. McA. Gordon, Knots, homology spheres, and contractible 4-manifolds, Topology, Vol. 14, No 2, 1975, 151-172.
- [K] R. C. Kirby, A calculus for framed links, Inventiones Math, 445, 1978, 35-56.
- [M] N. Martin, Some homology 3-spheres which bound acyclic 4-manifolds, Springer Lecture Notes, No 722, 1979, 85-92.
- [M1] N. Maruyama, Notes on homology 3-spheres which bound contractible 4-manifolds I. (to appear in Journal of Tsuda College, Vol 13)
- [M2] N. Maruyama, Notes on homology 3-spheres which bound contractible 4-manifolds II. (preprint)

NORIKO MARUYAMA

Department of Mathematics

Tsuda College

Kodaira, Tokyo, 187

Japan